

DOMAIN WALL PARTITION FUNCTIONS AND KP

O FODA, M WHEELER AND M ZUPARIC

ABSTRACT. We observe that the partition function of the six vertex model on a finite square lattice with domain wall boundary conditions is (a restriction of) a KP τ function and express it as an expectation value of charged free fermions (up to an overall normalization).

To Professor T Miwa on his 60th birthday.

0. INTRODUCTION

In [1], Korepin introduced domain wall boundary conditions for the six vertex model on an $N \times N$ square lattice, and obtained a set of conditions that determine the corresponding partition function Z_N . In [2], Izergin proposed a determinant expression for Z_N , and showed that it satisfies Korepin's conditions. In [3], Lascoux rewrote Z_N as a determinant of a product of two non-square matrices. In this note we observe that, given Lascoux's form, Z_N is (a restriction of) a KP τ function (for all values of the crossing parameter) and rewrite it as an expectation value of charged free fermions.

In section 1, we recall basic definitions related to the six vertex model and domain wall boundary conditions, including Korepin's conditions, and in section 2, we recall Izergin's determinant expression for Z_N , followed by Lascoux's expression, and observe that the latter is (a restriction of) a KP τ function. In section 3, we propose an expectation value of KP charged free fermions F_N^{free} that by construction is a KP τ function, and show that (under suitable restrictions of the KP time variables) F_N^{free} becomes equal to Z_N (up to an overall normalization). In section 4, we include a number of remarks.

1. DOMAIN WALL PARTITION FUNCTIONS

In this section, we list a number of basic definitions related to the six vertex model, domain wall boundary conditions, *etc.* For details, see [4, 5].

1.1. Oriented lattice lines and rapidity variables. Consider a square lattice with N horizontal lines (rows) and N vertical lines (columns) that intersect at N^2 points, and assign the i -th horizontal line an orientation from left to right and a rapidity x_i , and the j -th vertical line an orientation from bottom to top and a rapidity y_j , as in Figure 1.

1.2. Arrows, weights and a crossing parameter. Assign each line segment an arrow that can point in either direction, and define the vertex v_{ij} as the intersection point of the i -th horizontal line and the j -th vertical line, the four line segments attached to this intersection point and the arrows on these segments. Assign v_{ij} a weight w_{ij} that depends on the orientations of its arrows, the rapidities x_i and y_j and a *crossing parameter* μ that is the same for all vertices.

2000 *Mathematics Subject Classification.* Primary 82B20, 82B23.

Key words and phrases. Domain wall partition functions. Six vertex model. KP. Free fermions.

1.3. The six vertex model. Since an arrow on a line segment can point in either direction, there are $2^4 = 16$ possible (types of) vertices. In the six-vertex model, the weights of all vertices, except six, are zero [4]. The six vertices with non-zero weights form three pairs of equal-weight vertices. They are shown in Figure 2.




Figure 1. A square lattice with oriented lines and rapidity variables. The line orientations are indicated by white arrows.




Figure 2. The vertices of the six vertex model. Vertices that are in the same column share the weight shown below that column. The white arrows indicate the line orientations which are needed to completely specify the vertices.

In the notation of Figure 2, the weights of the six-vertex model are

$$(1) \quad w_{ij}^a = \sinh(-x_i + y_j + \mu), \quad w_{ij}^b = \sinh(-x_i + y_j), \quad w_{ij}^c = \sinh \mu$$

that satisfy the Yang-Baxter equations [4].

1.4. Domain wall boundary conditions. When all arrows on the left and right boundaries point inwards, and all arrows on the upper and lower boundaries point outwards, as in Figure 3, we obtain *domain wall boundary conditions*. The inner arrows remain free. Different inner arrow orientations lead to different *lattice configurations* that together constitute the statistical states of the model.

1.5. Domain wall partition functions. The weight of a lattice configuration is the product of the weights w_{ij} of its vertices v_{ij} . The weighted sum over all lattice configurations, on an $N \times N$ lattice with fixed domain wall boundary conditions, is a domain wall partition function Z_N .

$$(2) \quad Z_N \equiv Z_N(\mathbf{x}, \mathbf{y}, \mu) := \sum_{\substack{\text{all allowed} \\ \text{configurations}}} \prod_{\substack{\text{all} \\ \text{vertices}}} w_{ij}(x_i, y_j, \mu)$$




Figure 3. A lattice configuration with domain wall boundary conditions.

1.6. Korepin's conditions. In [1], Korepin obtained four conditions that uniquely determine $Z_N(\mathbf{x}, \mathbf{y}, \mu)$. These are

1. Z_N is a symmetric function in the $\{\mathbf{x}\}$ and in the $\{\mathbf{y}\}$ rapidity variables.
2. Z_N is a trigonometric polynomial of degree $(N - 1)$ in any rapidity variable.
3. Setting $x_1 = y_1 + \mu$, Z_N satisfies the recursion relation

$$Z_N|_{x_1=y_1+\mu} = \left(\prod_{i=2}^N \sinh(-x_i + y_1) \right) \sinh \mu \left(\prod_{j=2}^N \sinh(-x_1 + y_j) \right) Z_{(N-1)}(\hat{x}_1, \hat{y}_1, \mu)$$

where $\hat{x}_1, \hat{y}_1$ indicate that x_1 and y_1 are missing from the sets $\{\mathbf{x}\}$ and $\{\mathbf{y}\}$.

4. $Z_1 = \sinh \mu$

We refer to [1, 5] for details and proofs.

2. Z_N IS (A RESTRICTION OF) A KP τ FUNCTION

2.1. Izergin's determinant expression. Working in terms of the variables $u_i = e^{2x_i}$, $v_i = e^{2y_i}$, and $q = e^{-2\mu}$, Izergin's determinant expression is

$$(3) \quad Z_N(\mathbf{u}, \mathbf{v}, q) = c_N \frac{\prod_{i,j=1}^N (u_i - v_j)(qu_i - v_j)}{\prod_{1 \leq i < j \leq N} (u_i - u_j)(v_j - v_i)} \det \left[\frac{1}{(u_i - v_j)(qu_i - v_j)} \right]_{i,j=1}^N$$

where

$$(4) \quad c_N = (q-1)^N \prod_{i=1}^N u_i^{\frac{1}{2}} v_i^{\frac{1}{2}}$$

The virtue of Izergin's expression is that it is straightforward to show that it satisfies Korepin's conditions [2, 5].

2.2. Lascoux's determinant expression. In [3], Lascoux showed that Izergin's expression for Z_N can be rewritten as

$$(5) \quad Z_N(\mathbf{u}, \mathbf{v}, q) = c_N \det \left[\sum_{j=1}^{2N-1} \alpha_{i,j}(\mathbf{u}) \beta_{j,k}(\mathbf{v}, q) \right]_{i,k=1}^N$$

where

$$(6) \quad \alpha_{i,j}(\mathbf{u}) = h_{-i+j}(\mathbf{u}), \quad \beta_{j,k}(\mathbf{v}, q) = \frac{q^{j-k+1} - q^{k-1}}{q-1} e_{-j+k+N-1}(-\mathbf{v})$$

and $h_n(\mathbf{u})$ and $e_n(\mathbf{v})$ are the n -th homogeneous and the n -th elementary symmetric functions, respectively [6]. For a proof that Lascoux's expression is equivalent to Izergin's, see [3]. As we will see shortly, the virtue of Lascoux's expression is that it is straightforward to show that it is (a restriction of) a KP τ function.

2.3. Remarks on notation. In $\alpha_{i,j}$, $\beta_{i,j}$, and other matrix elements, i is the row and j is the column index. We will omit writing the dependence on the parameter q explicitly, as it is always there.

2.4. Partitions. We denote the partition of an integer $|\lambda|$ into parts $\lambda_1 \geq \lambda_2 \geq \dots$, by $[\lambda] = [\lambda_1, \lambda_2, \dots, \lambda_L]$, where L is the number of non-zero parts. m^n in $[\dots, m^n, \dots]$ indicates that $[\lambda]$ has n parts of length m . Further, m^0 indicates that there are no parts of length m and 0^n stands for n parts of length 0.

2.5. Young diagrams. Consider the Young diagram λ of a partition $[\lambda]$ as in Figure 4. The cells are labelled from top to bottom and from left to right. The cell with coordinates (i, j) , $i, j \geq 1$ is c_{ij} . The number of cells $c_{11}, c_{22}, \dots$, on the main diagonal is d .

(1,1)	(1,2)	(1,3)	(1,4)
(2,1)	(2,2)	(2,3)	
(3,1)	(3,2)	(3,3)	
(4,1)	(4,2)		

Figure 4. The Young diagram $\lambda = [4, 3, 3, 2] \equiv [4, 3^2, 2]$. The labels (i, j) in a cell c_{ij} are its coordinates. In this example, $d = 3$.

2.6. Hooks. Given a cell c_{ij} , we define a hook h_{ij} as the union of c_{ij} , the set of all cells below it and all cells to the right of it. We say that the hook h_{ij} has a corner at c_{ij} . The diagram on the left hand side of Figure 5 has 6 hooks, one for each cell. The right hand side of Figure 5 is the hook corresponding to the cell $c_{(2,1)}$.

2.7. Horizontal a -parts and vertical b -parts. We decompose a hook h_{ij} into a horizontal a -part a_{ij} of length $|a_{ij}|$ that consists of all cells to the right of c_{ij} , and a vertical b -part b_{ij} of length $|b_{ij}|$ that consists of c_{ij} and all cells below it. Hence the allowed a -part lengths are $0, 1, 2, \dots$, while the allowed b -part lengths are $1, 2, \dots$. The length of h_{ij} will be indicated by $|h_{ij}|$ as well, so that

$$(7) \quad |h_{ij}| = |a_{ij}| + |b_{ij}|$$

2.8. Frobenius coordinates of λ . Decomposing each hook h_{ii} , $i = 1, \dots, d$ that has a corner at a cell on the main diagonal of λ into the corner cell c_{ii} , the set of m_i cells to the right of c_{ii} , and n_i cells below it, we obtain $2d$ integers $\{m_i, n_j\}$ which define the Frobenius coordinates of λ .



Figure 5. On the left, the Young diagram corresponding to $\lambda = [2, 2, 2] \equiv [2^3]$. The labels $[a, b]$ in a cell are the a -part and b -part of the hook associated with that cell. On the right, the hook corresponding to c_{21} in λ . In this hook, the a -part a_{21} has length 1, the b -part b_{21} has length 2.

2.9. Expanding Z_N in terms of Schur functions. To make contact with KP theory and eventually obtain an expression for Z_N as a vacuum expectation value of free fermions, we start by expanding Lascoux's expression in terms of Schur functions. Using (6) and the Cauchy-Binet identity in (5), we obtain

$$(8) \quad Z_N(\mathbf{u}, \mathbf{v}) = c_N \sum_{1 \leq j_1 < \dots < j_N \leq 2N-1} \det [h_{-i+j_l}(\mathbf{u})]_{i,l=1}^N \det [\beta_{j_l, k}(\mathbf{v})]_{l,k=1}^N$$

Rewriting j_l as $j_l = \lambda_{(N+1)-l} + l$, then reversing the order of all rows and all columns in the first matrix and transposing it, the sum on the right hand side of (8) becomes

$$(9) \quad \sum_{0 \leq \lambda_N \leq \dots \leq \lambda_1 \leq N-1} \det [h_{\lambda_i - i + l}(\mathbf{u})]_{i,l=1}^N \det [\beta_{\lambda_{(N+1)-l} + l, k}(\mathbf{v})]_{l,k=1}^N$$

Finally, using the definition of Schur functions

$$(10) \quad S_\lambda(\mathbf{u}) = \det [h_{\lambda_i - i + l}(\mathbf{u})]_{i,l=1}^N$$

and (6) to write

$$(11) \quad \begin{aligned} c_\lambda^{(N)}(\mathbf{v}) &= \det \left[\beta_{(\lambda_{l_{conj}}) - l_{conj} + (N+1), k}(\mathbf{v}) \right]_{l,k=1}^N \\ &= \det \left[\frac{q^{(\lambda_{l_{conj}}) - l_{conj} - k + N + 2} - q^{k-1}}{q-1} e_{-(\lambda_{l_{conj}}) + l_{conj} + k - 2}(-\mathbf{v}) \right]_{l,k=1}^N \end{aligned}$$

where

$$(12) \quad l_{conj} = (N+1) - l$$

we end up with

$$(13) \quad Z_N(\mathbf{u}, \mathbf{v}) = c_N \sum_{\lambda \subseteq [(N-1)^N]} c_\lambda^{(N)}(\mathbf{v}) S_\lambda(\mathbf{u})$$

Since the summation in (13) is over partitions of at most N parts, and there are N independent u -variables, and N independent v -variables, there are no terms in the expansion that trivially vanish due to lack of sufficiently many independent variables.

2.10. Expanding Z_N in terms of character polynomials. Next, following [7], we introduce the power sum variables

$$(14) \quad t_n = \frac{1}{n} \sum_{i=1}^N u_i^n, \quad n = 1, 2, \dots$$

and the polynomials $p_n(\mathbf{t})$

$$(15) \quad p_n(\mathbf{t}) = \sum_{n_1+2n_2+3n_3+\dots=n} \frac{t_1^{n_1} t_2^{n_2} t_3^{n_3} \dots}{n_1! n_2! n_3! \dots}$$

where $p_i(\mathbf{t}) = 0$ for $i < 0$. Next, we define the polynomials $h_{m,n}(\mathbf{t})$

$$(16) \quad h_{m,n}(\mathbf{t}) := (-1)^n \sum_{k=0}^n p_{k+m+1}(\mathbf{t}) p_{n-k}(-\mathbf{t})$$

and finally the character polynomials $\chi_\lambda(\mathbf{t})$

$$(17) \quad \chi_\lambda(\mathbf{t}) := \det [h_{m_i, n_j}(\mathbf{t})]_{i,j=1}^d$$

where $\{m_i, n_j\}$ are the Frobenius coordinates of λ , and the upper limit d is the number of cells on the main diagonal. In terms of character polynomials and the t -variables of (14) (and by slightly abusing notation), we rewrite $Z_N(\mathbf{u}, \mathbf{v})$ as

$$(18) \quad Z_N(\mathbf{t}, \mathbf{v}) = c_N \sum_{\lambda \subseteq (N-1)^N} c_\lambda^{(N)}(\mathbf{v}) \chi_\lambda(\mathbf{t})$$

where $c_\lambda^{(N)}(\mathbf{v})$ was defined in (11). We conclude that Z_N can be expanded in terms of character polynomials, with coefficients that are minors of a determinant such that (by construction) they satisfy Plücker relations [7]. If the t -variables were all independent, then $Z_N(\mathbf{t}, \mathbf{v})$ would be identically a KP τ function. We refer to [7] for details and proofs.

2.11. Restricted t -variables. $Z_N(\mathbf{u}, \mathbf{v})$ is a function of N independent u -variables. On the other hand, from the definition of the character polynomials, it is easy to show that $Z_N(\mathbf{t}, \mathbf{v})$ requires in general more than N t -variables. Since the latter are power sums of the N u -variables, they cannot be all independent. Let us write $Z_N^{rest}(\mathbf{t}, \mathbf{v})$ to emphasise the fact that the t -variables are restricted to be power sums of a (generally) smaller number of u -variables. This restriction spoils the interpretation of the t -variables as KP time variables, and of $Z_N^{rest}(\mathbf{t}, \mathbf{v})$ as a KP τ function.

2.12. Free t -variables. We can formally consider $Z_N^{rest}(\mathbf{t}, \mathbf{v})$ as our starting point, take the t -variables to be independent, write it as $Z_N^{free}(\mathbf{t}, \mathbf{v})$ and think of $Z_N^{rest}(\mathbf{t}, \mathbf{v})$ as a restriction of $Z_N^{free}(\mathbf{t}, \mathbf{v})$ obtained by setting the t -variables to power sums of the N u -variables using (14). Viewed this way, $Z_N^{free}(\mathbf{t}, \mathbf{v})$ is a KP τ function, and $Z_N^{rest}(\mathbf{t}, \mathbf{v}) = Z_N(\mathbf{u}, \mathbf{v})$, is a restriction of a KP τ function.

3. FERMIONIC EXPRESSION FOR Z_N

Given that any KP τ function can be written as the vacuum expectation value of exponentials in neutral bilinears in KP charged fermions¹ [7], we wish to write Z_N^{free} as such. Our plan is to propose a vacuum expectation value $F_N^{free}(\mathbf{t}, \mathbf{v})$, where we use the superscript *free* to emphasise that the t -variables are independent, that by construction is a KP τ function, then show that $F_N^{free}(\mathbf{t}, \mathbf{v}) = Z_N^{free}(\mathbf{t}, \mathbf{v})$. Hence restricting the t -variables in $F_N^{free}(\mathbf{t}, \mathbf{v})$ to be power sums of N u -variables, one recovers $Z_N^{rest}(\mathbf{t}, \mathbf{v})$, and $Z_N(\mathbf{u}, \mathbf{v})$, as an expectation value of charged free fermions.

3.1. Charged free fermions. Consider the free fermion operators $\{\psi_n, \psi_n^*\}$, $n \in \mathbb{Z}$, with charges $\{+1, -1\}$ and energies n . They generate a Clifford algebra over $\mathbb{C}$ defined by the anti-commutation relations

$$(19) \quad \left. \begin{aligned} [\psi_m, \psi_n]_+ &= 0 \\ [\psi_m^*, \psi_n^*]_+ &= 0 \\ [\psi_m, \psi_n^*]_+ &= \delta_{m,n} \end{aligned} \right\} \quad \forall m, n \in \mathbb{Z}$$

3.2. Vacuum states. We define the vacuum states $\langle 0|$ and $|0\rangle$ by the actions

$$(20) \quad \left. \begin{aligned} \langle 0|\psi_n = \psi_m|0\rangle &= 0, \\ \langle 0|\psi_m^* = \psi_n^*|0\rangle &= 0, \end{aligned} \right\} \quad \forall m < 0, n \geq 0$$

and we adopt the inner product normalization

$$(21) \quad \langle 0|0\rangle = 1$$

3.3. Creation and annihilation operators. We refer to the operators that annihilate the vacuum state $|0\rangle$, $\psi_m|0\rangle = 0$, $m = -1, -2, \dots$, and $\psi_n^*|0\rangle = 0$, $n = 0, 1, 2, \dots$ as *annihilation operators*, and to the rest, $\psi_m|0\rangle \neq 0$, $m = 0, 1, 2, \dots$, and $\psi_n^*|0\rangle \neq 0$, $n = -1, -2, \dots$ as *creation operators*.

3.4. Normal ordering. The normal-ordered product is defined, as usual, by placing annihilation operators to the right of creation operators

$$: \psi_i \psi_j^* : = \psi_i \psi_j^* - \langle 0|\psi_i \psi_j^*|0\rangle$$

3.5. The Heisenberg algebra. Consider the operators H_m ,

$$(22) \quad H_m := \sum_{j \in \mathbb{Z}} : \psi_j \psi_{j+m}^* :, \quad m \in \mathbb{Z}$$

that together with the central element 1 form a Heisenberg algebra

$$(23) \quad [H_m, H_n] = m \delta_{m+n, 0}, \quad \forall m, n \in \mathbb{Z}$$

¹By this, we mean linear combinations of terms of the form $\psi_m \psi_n^*$.

3.6. The Hamiltonian. Using $H_m, m = 1, 2, 3, \dots$, we define the Hamiltonian

$$(24) \quad H(\mathbf{t}) := \sum_{m=1}^{\infty} t_m H_m$$

3.7. Fermion bilinears. Given a partition λ , we can associate to each hook h_{ij} a bilinear $\psi_{|a_{ij}|} \psi_{-|b_{ij}|}^*$ with net charge zero. Both fermions in the bilinear are creation operators,

$$(25) \quad \psi_{|a_{ij}|} \psi_{-|b_{ij}|}^* = -\psi_{-|b_{ij}|}^* \psi_{|a_{ij}|}$$

and normal ordering is unnecessary.

3.8. Boson-fermion correspondence. It is possible to realise fermionic Fock space expressions as elements in the polynomial ring $\mathbb{C}[t_1, t_2, \dots]$ [7]. For the purposes of this work, all we need is that fact that the character polynomial $\chi_\lambda(\mathbf{t})$ can be generated as follows

$$(26) \quad \langle 0 | e^{H(\mathbf{t})} \psi_{-b_1}^* \cdots \psi_{-b_d}^* \psi_{a_d} \cdots \psi_{a_1} | 0 \rangle = (-1)^{b_1 + \cdots + b_d} \chi_\lambda(\mathbf{t})$$

where $a_d < \cdots < a_1$ and $b_d < \cdots < b_1$ are the a -part and b -part lengths of the hooks associated with cells $c_{11}, c_{22}, \dots$, on the main diagonal of λ , and d is the number of these cells. In terms of Frobenius coordinates, $a_i = m_i$ and $b_j = n_j + 1$.

3.9. Partitions and fermion monomials. Consider the fermion monomial consisting of d ψ and d ψ^* fermions, $\psi_{-b_1}^* \cdots \psi_{-b_d}^* \psi_{a_d} \cdots \psi_{a_1}$, where $a_d < \cdots < a_1$ and $b_d < \cdots < b_1$ as on the left hand side of (26). The fermions in the monomial are ordered according to length and correspond to the a -parts and b -parts of the partition λ on the right hand side of (26), with the longer b -parts to the left and the longer a -parts to the right. We refer to such a monomial associated to a partition λ as a partition-ordered, or λ -ordered monomial. The partition in Figure 4 has $d = 3$, a set of b -parts $\{4, 3, 1\}$ and a set of a -parts $\{3, 1, 0\}$. The corresponding monomial is $\psi_{-4}^* \psi_{-3}^* \psi_{-1}^* \psi_0 \psi_1 \psi_3$.

3.10. A fermion expectation value. Consider the vacuum expectation value

$$(27) \quad F_N^{\text{free}}(\mathbf{t}, \mathbf{v}) = \langle 0 | e^{H(\mathbf{t})} e^{X_0^{(N)}} e^{X_1^{(N)}} \cdots e^{X_{N-2}^{(N)}} | 0 \rangle$$

where

$$(28) \quad X_a^{(N)} = \sum_{b=1}^N (-)^b d_{[a+1, 1^{b-1}]}^{(N)} \psi_{-b}^* \psi_a, \quad a = 0, 1, \dots, N-2$$

where 1^0 inside a partition $[\lambda]$ stands for a trivial, length-0 part, and

$$(29) \quad d_\lambda^{(N)} = c_\lambda^{(N)} / c_\phi^{(N)}$$

The coefficient $c_\lambda^{(N)}$ was defined in (11), and the (infinitely many) t -parameters in (27) are taken as independent variables rather than power sums of the (finitely many) u -parameters. Since F_N^{free} is an expectation value of exponentials of neutral bilinears in $\{\psi_n, \psi_m^*\}$, which are generators of $gl(\infty)$, followed by the time evolution operator $e^{H(\mathbf{t})}$, it is by construction a KP τ function [7]. Since all fermions in the linear combinations of bilinears in (28) are creation operators, there is no need for normal ordering these terms.

3.11. A combinatorial interpretation. The exponentials that appear in F_N^{free} in (27) have the following interpretation. Consider a rectangular partition of N rows and $(N-1)$ columns. For example, for $N=3$, see the diagram on the left hand side of Figure 5. For the j -th column, with N cells of a -part length $a = (N-1)-j$, we have an exponential $X_a^{(N)}$, $0 \leq a \leq (N-2)$, which is a sum of N terms. The b -th term is of the form $(-1)^b d_{[h]}^{(N)} \psi_{-b}^* \psi_a$, where $[h]$ is the partition corresponding to the hook that has a corner at that cell, and a and b are the a -part and b -part lengths of that hook.

3.12. F_N is proportional to Z_N . In the rest of this work, we show that

$$(30) \quad c_N c_\phi^{(N)}(\mathbf{v}) F_N^{free}(\mathbf{t}, \mathbf{v}) = Z_N^{free}(\mathbf{t}, \mathbf{v})$$

3.13. The normalization factor. Expanding the exponentials in (27) and using $\langle 0|0 \rangle = 1$, the character polynomial expansion of F_N^{free} starts with 1. On the other hand, since $\chi_\phi(\mathbf{t}) = 1$, the character polynomial expansion of Z_N in (18) starts with the t -independent term, $c_N c_\phi^{(N)}(\mathbf{v})$. We choose to keep the definition of the vertex weights as in (1), and show that F_N and Z_N are equal up to a factor of $c_N c_\phi^{(N)}(\mathbf{v})$.

3.14. Example: $N=3$. Before we prove (30) for general N , we verify it in the simple but non-trivial case of $N=3$. Omitting the superscripts (3) to simplify the notation, we have

$$(31) \quad \begin{aligned} c_\lambda &= \det \left[\frac{q^{\binom{\lambda_{4-i}}{2} + i - j + 1} - q^{j-1}}{q-1} e^{-\binom{\lambda_{4-i}}{2} - i + j + 2} (-\mathbf{v}) \right]_{i,j=1}^3 \\ X_0 &= -d_{[1,0,0]} \psi_{-1}^* \psi_0 + d_{[1,1,0]} \psi_{-2}^* \psi_0 - d_{[1,1,1]} \psi_{-3}^* \psi_0 \\ X_1 &= -d_{[2,0,0]} \psi_{-1}^* \psi_1 + d_{[2,1,0]} \psi_{-2}^* \psi_1 - d_{[2,1,1]} \psi_{-3}^* \psi_1 \end{aligned}$$

where d_λ was defined in (29). Notice that in this case, the relevant partition is shown on the left hand side of Figure 5, and the bilinears in X_0 and X_1 correspond to the hooks of the cells in the two columns. This leads to

$$(32) \quad \begin{aligned} e^{X_0} e^{X_1} &= 1 - d_{[1,0,0]} \psi_{-1}^* \psi_0 + d_{[1,1,0]} \psi_{-2}^* \psi_0 - d_{[1,1,1]} \psi_{-3}^* \psi_0 \\ &\quad - d_{[2,0,0]} \psi_{-1}^* \psi_1 + d_{[2,1,0]} \psi_{-2}^* \psi_1 - d_{[2,1,1]} \psi_{-3}^* \psi_1 \\ &\quad + \left(d_{[1,1,0]} d_{[2,0,0]} - d_{[1,0,0]} d_{[2,1,0]} \right) \psi_{-2}^* \psi_{-1}^* \psi_0 \psi_1 \\ &\quad + \left(d_{[1,0,0]} d_{[2,1,1]} - d_{[1,1,1]} d_{[2,0,0]} \right) \psi_{-3}^* \psi_{-1}^* \psi_0 \psi_1 \\ &\quad + \left(d_{[1,1,1]} d_{[2,1,0]} - d_{[1,1,0]} d_{[2,1,1]} \right) \psi_{-3}^* \psi_{-2}^* \psi_0 \psi_1 \end{aligned}$$

Consider the six fermion bilinears on the right hand side of (32). Using the boson-fermion correspondence, each of these bilinears corresponds to a partition $[\lambda]$ (which in the case of bilinears is a single hook), and its coefficient $d_{[\lambda]}$ corresponds to the same partition $[\lambda]$. If the same is true for the three fermion quartic terms, then from (18), F_3 would be proportional to Z_3 . To put the coefficients of the quartic terms in the right form, we need to show that

$$(33) \quad d_{[1,1,0]}d_{[2,0,0]} - d_{[1,0,0]}d_{[2,1,0]} = -d_{[2,2,0]}$$

$$(34) \quad d_{[1,0,0]}d_{[2,1,1]} - d_{[1,1,1]}d_{[2,0,0]} = d_{[2,2,1]}$$

$$(35) \quad d_{[1,1,1]}d_{[2,1,0]} - d_{[1,1,0]}d_{[2,1,1]} = -d_{[2,2,2]}$$

3.15. Plücker relations. Equations (33–35) are examples of Plücker relations, which are identities involving bilinears in $n \times n$ minors of a $2n \times 2n$ matrix with zero determinant. In this case, $n = 3$. For details, see [8]. Using the notation

$$(36) \quad c_\lambda \equiv c_{[\lambda_1, \lambda_2, \lambda_3]} = \det [\gamma_{\lambda_3+1}, \gamma_{\lambda_2+2}, \gamma_{\lambda_1+3}] = |\gamma_{\lambda_3+1}, \gamma_{\lambda_2+2}, \gamma_{\lambda_1+3}|,$$

where γ_b is the 3-component column vector

$$(37) \quad \gamma_b = \left(\frac{q^{b-a+1} - q^{a-1}}{q-1} e_{-b+a+2}(-\mathbf{v}) \right)_{a=1,2,3},$$

we consider the 6×6 determinant

$$(38) \quad \begin{vmatrix} \gamma_{\mu_1} & \gamma_{\mu_2} & \gamma_{\nu_1} & \gamma_{\nu_2} & \gamma_{\nu_3} & \gamma_{\nu_4} \\ 0 & 0 & \gamma_{\nu_1} & \gamma_{\nu_2} & \gamma_{\nu_3} & \gamma_{\nu_4} \end{vmatrix} = 0,$$

and Laplace expand it as the sum of bilinears in 3×3 determinants

$$(39) \quad \begin{aligned} & |\gamma_{\mu_1}, \gamma_{\mu_2}, \gamma_{\nu_1}| |\gamma_{\nu_2}, \gamma_{\nu_3}, \gamma_{\nu_4}| - |\gamma_{\mu_1}, \gamma_{\mu_2}, \gamma_{\nu_2}| |\gamma_{\nu_1}, \gamma_{\nu_3}, \gamma_{\nu_4}| \\ & + |\gamma_{\mu_1}, \gamma_{\mu_2}, \gamma_{\nu_3}| |\gamma_{\nu_1}, \gamma_{\nu_2}, \gamma_{\nu_4}| - |\gamma_{\mu_1}, \gamma_{\mu_2}, \gamma_{\nu_4}| |\gamma_{\nu_1}, \gamma_{\nu_2}, \gamma_{\nu_3}| = 0 \end{aligned}$$

We can use (39) to generate the Plücker relations that we need by suitably choosing the column vectors γ_b .

3.16. The $N = 3$ Plücker relations. The required Plücker relation, defined as in (41), has a term $d_{[\lambda]}d_{[\phi]}$, where $d_{[\phi]} = 1$, and $\lambda = [2, 2, 0]$. From (36) we see that the bilinear that we need is $|\gamma_{1+0}, \gamma_{2+2}, \gamma_{3+2}| |\gamma_{1+0}, \gamma_{2+0}, \gamma_{3+0}| = |\gamma_1, \gamma_4, \gamma_5| |\gamma_1, \gamma_2, \gamma_3|$. This dictates the choice of parameters

$$(40) \quad (\mu_1, \mu_2, \nu_1, \nu_2, \nu_3, \nu_4) = (1, 4, 5, 1, 2, 3)$$

Using (40) in (39), and the antisymmetry property of determinants, we obtain

$$(41) \quad |\gamma_1, \gamma_4, \gamma_5| |\gamma_1, \gamma_2, \gamma_3| - |\gamma_1, \gamma_3, \gamma_5| |\gamma_1, \gamma_2, \gamma_4| = -|\gamma_1, \gamma_3, \gamma_4| |\gamma_1, \gamma_2, \gamma_5|$$

which, using (29) and (36), is the Plücker relation that we need to prove (33).

Equations (34–35) are obtained by setting $(\mu_1, \mu_2, \nu_1, \nu_2, \nu_3, \nu_4)$ to $(2, 4, 5, 1, 2, 3)$, then to $(3, 4, 5, 1, 2, 3)$ in (39). Finally, using (26), we obtain

$$(42) \quad c_\phi \langle 0 | e^{H(\mathbf{t})} e^{X_0} e^{X_1} | 0 \rangle = \sum_{\lambda \subseteq [2,2,2]} c_\lambda(\mathbf{v}) \chi_\lambda(\mathbf{t})$$

The right hand side of (42) is Lascoux's form expanded as in (18). The above example shows that the proof of (30) relies on using Plücker relations to simplify the coefficients of expansions of the exponentials that appear in (27).

3.17. Plan of proof of Equation (30). Firstly, we expand the product of two exponentials in terms of neutral monomials in the fermions, with coefficients that can be simplified using Plücker relations, then we use induction to show that this can be done for the product of N exponentials. Finally we use boson-fermion correspondence to obtain Lascoux's form of Z_N .

3.18. Remark on notation. In the following, we consider only the general N case, so we omit the superscripts (N) , and write X_a , instead of $X_a^{(N)}$, and so on.

3.19. Expanding two exponentials. For $0 \leq a_2 < a_1 \leq (N-2)$, consider the product

$$(43) \quad e^{X_{a_2}} e^{X_{a_1}} = 1 + X_{a_2} + X_{a_1} + X_{a_2} X_{a_1}$$

where

$$(44) \quad \begin{aligned} X_{a_1} &= \sum_{b=1}^N (-1)^b d_{[a_1+1, 1^{(b-1)}]} \psi_{-b}^* \psi_{a_1} \\ X_{a_2} &= \sum_{b=1}^N (-1)^b d_{[a_2+1, 1^{(b-1)}]} \psi_{-b}^* \psi_{a_2} \\ X_{a_2} X_{a_1} &= \sum_{\substack{b_1, b_2=1 \\ b_1 \neq b_2}}^N (-1)^{b_1+b_2} d_{[a_2+1, 1^{(b_2-1)}]} d_{[a_1+1, 1^{(b_1-1)}]} \psi_{-b_2}^* \psi_{a_2} \psi_{-b_1}^* \psi_{a_1} \end{aligned}$$

Note that while we know that $a_2 < a_1$, we can have $b_2 < b_1$ and $b_1 < b_2$. We account for these two possibilities by rewriting the cross-term as

$$(45) \quad X_{a_2} X_{a_1} = \left(\sum_{1 \leq b_2 < b_1 \leq N} + \sum_{1 \leq b_1 < b_2 \leq N} \right) (-1)^{b_1+b_2} d_{[a_2+1, 1^{(b_2-1)}]} d_{[a_1+1, 1^{(b_1-1)}]} \times \psi_{-b_1}^* \psi_{-b_2}^* \psi_{a_2} \psi_{a_1}$$

Relabelling the indices in the second summation, so that $b_1 \rightarrow b_2$ and $b_2 \rightarrow b_1$, and using the anticommutation relations to put the product of fermion operators in the form used in (26), the right hand side of (45) becomes

$$(46) \quad \sum_{1 \leq b_2 < b_1 \leq N} (-1)^{b_1+b_2} \left(d_{[a_2+1, 1^{(b_2-1)}]} d_{[a_1+1, 1^{(b_1-1)}]} - d_{[a_2+1, 1^{(b_1-1)}]} d_{[a_1+1, 1^{(b_2-1)}]} \right) \times \psi_{-b_1}^* \psi_{-b_2}^* \psi_{a_2} \psi_{a_1}$$

we see that once again we need to simplify the coefficients. Following the same ideas that were used in the $N=3$ example above, we write

$$(47) \quad c_\lambda = \left| \gamma_{(\lambda_N)+1}, \gamma_{(\lambda_{N-1})+2}, \dots, \gamma_{(\lambda_1)+N} \right|$$

where γ_b is the N -component column vector

$$(48) \quad \gamma_b = \left(\frac{q^{b-a+1} - q^{a-1}}{q-1} e_{-b+a+N-1}(-\mathbf{v}) \right)_{a=1, \dots, N}$$

then consider the $2N \times 2N$ determinant expression

$$(49) \quad \begin{vmatrix} \gamma_{\mu_1} & \cdots & \gamma_{\mu_{N-1}} & \gamma_{\nu_1} & \gamma_{\nu_2} & \cdots & \gamma_{\nu_{N+1}} \\ 0 & \cdots & 0 & \gamma_{\nu_1} & \gamma_{\nu_2} & \cdots & \gamma_{\nu_{N+1}} \end{vmatrix} = 0$$

and Laplace expand it as a sum of bilinears in $N \times N$ determinants

$$(50) \quad \sum_{p=1}^{N+1} (-1)^{p+1} |\gamma_{\mu_1} \cdots \gamma_{\mu_{N-1}} \gamma_{\nu_p}| |\gamma_{\nu_1} \cdots \hat{\gamma}_{\nu_p} \cdots \gamma_{\nu_{N+1}}| = 0$$

where $\hat{\gamma}$ indicates a missing term.

3.20. The general N Plücker relation. The procedure outlined for the $N = 3$ case in subsection **3.16**, can be extended to general N . From (46), the Plücker relation that we need contains a bilinear term $d_{[\lambda]} d_{[\phi]}$, where λ has two a -parts, $a_1 > a_2$ and two b -parts, $b_1 > b_2$, so that

$$(51) \quad \lambda = [a_1 + 1, a_2 + 2, 2^{b_2-2}, 1^{b_1-b_2-1}, 0^{N-b_1}],$$

where we have indicated the 0-length elements as well to make the total length of the partition N . We prepare a $2N$ -element sequence Σ^0

$$(52) \quad \Sigma^0 = [1, 2, \dots, N | 1, 2, \dots, N]$$

where the vertical bar $|$ in (52) emphasizes the structure of Σ^0 as a concatenation of a left sequence $\Sigma_{left}^0 = [1, 2, \dots, N]$ and a right sequence $\Sigma_{right}^0 = [1, 2, \dots, N]$. Given $\lambda = [\lambda_1, \dots, \lambda_N]$, where $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_N$, we modify Σ_{left}^0 as follows. We add λ_1 to the largest integer (N) in Σ_{left}^0 , λ_2 to the next largest ($N-1$), and so forth. Σ_{right}^0 remains unmodified. This generates a new sequence Σ . Choosing the parameters in the $2N$ -element sequence $(\mu_1, \dots, \mu_{N-1}, \nu_1, \dots, \nu_{N+1})$ by identifying them with the corresponding integers in Σ , we end up with

$$(53) \quad \begin{aligned} (\mu_1, \dots, \mu_{N-b_1}) &= (1, \dots, N-b_1) \\ (\mu_{N-b_1+1}, \dots, \mu_{N-b_2-1}) &= (N-b_1+2, \dots, N-b_2) \\ (\mu_{N-b_2}, \dots, \mu_{N-2}) &= (N-b_2+2, \dots, N) \\ (\mu_{N-1}, \nu_1) &= (N+a_2+1, N+a_1+1) \\ (\nu_2, \dots, \nu_{N+1}) &= (1, \dots, N) \end{aligned}$$

where ‘ $\dots$ ’ between two integers m and n stands for the integers $m+1, m+2, \dots, n-1$, that increase by 1 at a time. Using (50), the above choice of parameters leads to the Plücker relation

$$(54) \quad \begin{aligned} &|\Sigma, \gamma_{N+a_1+1}| |\gamma_1, \dots, \gamma_N| \\ &+ (-1)^{N-b_1} |\Sigma, \gamma_{N-b_1+1}| |\gamma_{N+a_1+1}, \gamma_1, \dots, \hat{\gamma}_{N-b_1+1}, \dots, \gamma_N| \\ &+ (-1)^{N-b_2} |\Sigma, \gamma_{N-b_2+1}| |\gamma_{N+a_1+1}, \gamma_1, \dots, \hat{\gamma}_{N-b_2+1}, \dots, \gamma_N| \\ &= 0 \end{aligned}$$

where Σ in (54) is the sequence

$$(55) \quad \Sigma = \{\gamma_1, \dots, \gamma_{N-b_1}, \gamma_{N-b_1+2}, \dots, \gamma_{N-b_2}, \gamma_{N-b_2+2}, \dots, \gamma_N, \gamma_{N+a_2+1}\}$$

Using (36), (54) can be written as

$$(56) \quad c_{[a_2+1, 1^{b_2-1}]} c_{[a_1+1, 1^{b_1-1}]} - c_{[a_2+1, 1^{b_1-1}]} c_{[a_1+1, 1^{b_2-1}]} = c_\phi c_{[a_1+1, a_2+2, 2^{b_2-1}, 1^{b_1-b_2-1}]}$$

Substituting (56) in the second line of (45), we obtain

$$(57) \quad X_{a_2} X_{a_1} = \sum_{1 \leq b_2 < b_1 \leq N} (-1)^{b_2+b_1} d_{[a_1+1, a_2+2, 2^{b_2-1}, 1^{b_1-b_2-1}]} \psi_{-b_1}^* \psi_{-b_2}^* \psi_{a_2} \psi_{a_1}$$

3.21. Expanding more than two exponentials. We wish to show that for all $0 \leq a_k < \dots < a_1 \leq N-2$, we have

$$(58) \quad X_{a_k} \dots X_{a_1} = \sum_{1 \leq b_k < \dots < b_1 \leq N} (-1)^{b_1+\dots+b_k} d_{\lambda_{(d=k)}} \psi_{-b_1}^* \dots \psi_{-b_k}^* \psi_{a_k} \dots \psi_{a_1}$$

where

$$(59) \quad \lambda_{(d=k)} = [a_1+1, a_2+2, \dots, a_k+k, k^{b_k-1}, (k-1)^{b_{k-1}-b_k-1}, \dots, 1^{b_1-b_2-1}, 0^{N-b_1}]$$

is the partition, with k cells on the main diagonal, associated with the fermion monomial on the right hand side of (58).

We start by assuming that (58) holds for some value of k , and show that it also holds for $k+1$. Since it holds for $k=2$, it holds for all k . Consider the expression

$$(60) \quad X_{a_k} \dots X_{a_1} X_{a_0} = \sum_{1 \leq b_k < \dots < b_1 \leq N} \sum_{\substack{b_0=1 \\ b_0 \neq b_1, \dots, b_k}}^N (-1)^{b_0+b_1+\dots+b_k} d_{\lambda_{(d=k)}} d_{[a_0+1, 1^{b_0-1}]} \psi_{-b_1}^* \dots \psi_{-b_k}^* \psi_{a_k} \dots \psi_{a_1} \psi_{-b_0}^* \psi_{a_0}$$

for $0 \leq a_k < \dots < a_1 < a_0 \leq (N-2)$, where $\lambda_{(d=k)}$ is still the same partition as in (58), and rewrite the summation as

$$(61) \quad \sum_{1 \leq b_k < \dots < b_1 \leq N} \sum_{\substack{b_0=1 \\ b_0 \neq b_1, \dots, b_k}}^N = \sum_{1 \leq b_k < \dots < b_0 \leq N} + \sum_{1 \leq b_k < \dots < b_2 < b_0 < b_1 \leq N} + \dots + \sum_{1 \leq b_k < b_0 < b_{k-1} < \dots < b_1 \leq N} + \sum_{1 \leq b_0 < b_k < \dots < b_1 \leq N}$$

Ordering the fermions in each summation, so that they are in the same order as in (26), and relabeling the indices, we obtain

$$(62) \quad \sum_{1 \leq b_k < \dots < b_0 \leq N} (-1)^{b_0+\dots+b_k} \left(\sum_{p=0}^k (-1)^p d_{\lambda_{(d=k|p)}} d_{[a_0+1, 1^{(b_p-1)}]} \right) \times \psi_{-b_0}^* \dots \psi_{-b_k}^* \psi_{a_k} \dots \psi_{a_0}$$

where $\lambda_{(d=k|p)}$ stands for $\lambda_{(d=k)}$, but the b -part that used to be b_p is now omitted and the b -part b_0 is added in correct position. For example, if $\lambda_{(d=6)}$ has b -parts $\{b_6 < b_5 < b_4 < b_3 < b_2 < b_1\}$, then $\lambda_{(d=6|3)}$ has $\{b_6 < b_5 < b_4 < b_2 < b_1 < b_0\}$. $\lambda_{(d=k|0)}$ indicates that no changes were made.

To simplify the sum in large brackets in (62), we consider (50), and, following the same procedure as before, use the evaluations

$$\begin{aligned}
(63) \quad & (\mu_1, \dots, \mu_{N-b_0}) = (1, \dots, N-b_0) \\
& (\mu_{N-b_0+1}, \dots, \mu_{N-b_1-1}) = (N-b_0+2, \dots, N-b_1) \\
& (\mu_{N-b_1}, \dots, \mu_{N-b_2-2}) = (N-b_1+2, \dots, N-b_2) \\
& (\mu_{N-b_2-1}, \dots, \mu_{N-b_3-3}) = (N-b_2+2, \dots, N-b_3) \\
& \vdots \\
& (\mu_{N-b_k-(k-1)}, \dots, \mu_{N-(k+1)}) = (N-b_k+2, \dots, N) \\
& (\mu_{N-k}, \dots, \mu_{N-1}, \nu_1) = (N+a_k+1, \dots, N+a_1+1, N+a_0+1) \\
& (\nu_2, \dots, \nu_{N+1}) = (1, \dots, N)
\end{aligned}$$

The required Plücker relation is

$$\begin{aligned}
(64) \quad & |\Sigma, \gamma_{N+a_0+1}| |\gamma_1, \dots, \gamma_N| \\
& + (-1)^{N-b_0+1} |\Sigma, \gamma_{N-b_0+1}| |\gamma_{N+a_0+1}, \dots, \hat{\gamma}_{N-b_0+1}, \dots, \gamma_N| \\
& + (-1)^{N-b_1+1} |\Sigma, \gamma_{N-b_1+1}| |\gamma_{N+a_0+1}, \dots, \hat{\gamma}_{N-b_1+1}, \dots, \gamma_N| \\
& + \dots \\
& + (-1)^{N-b_k+1} |\Sigma, \gamma_{N-b_k+1}| |\gamma_{N+a_0+1}, \dots, \hat{\gamma}_{N-b_k+1}, \dots, \gamma_N| \\
& = 0
\end{aligned}$$

where the sequence Σ in (64) is obtained as follows. Consider the sets of indices in (63) from top to bottom except the very last, $(1, \dots, N)$, and order them from left to right, so that the top-most $(1, \dots, N-b_0)$ becomes the left-most and $(N+a_k+1, \dots, N+a_1+1, N+a_0+1)$ becomes the right-most. Concatenate them and remove the right-most element $(N+a_0+1)$. Use the resulting $(N-1)$ -element sequence in the same order to generate the indices of Σ in (64). Using (36), (64) can be rewritten as

$$(65) \quad \sum_{p=0}^k (-1)^p c_{\lambda_{(d=k|p)}} c_{[a_0+1, 1^{b_p-1}]} = c_\phi c_{\lambda_{(d=k+1)}}$$

where $\lambda_{(d=k+1)}$ is the partition with $(k+1)$ cells on the main diagonal

$$(66) \quad \lambda_{(d=k+1)} = [a_0+1, a_1+2, \dots, a_k+(k+1), (k+1)^{b_k-1}, k^{b_{k-1}-b_k-1}, \dots, 1^{b_0-b_1-1}]$$

Relabeling a_i and b_j so that the indices take values from 1 to $(k+1)$, rather than from 0 to k , such that $a_1 > a_2 > \dots > a_{k+1}$ and $b_1 > b_2 > \dots > b_{k+1}$, we obtain

$$(67) \quad X_{a_{k+1}} \dots X_{a_1} = \sum_{1 \leq b_{k+1} < \dots < b_1 \leq N} (-1)^{b_1 + \dots + b_{k+1}} d_{\lambda_{(d=k+1)}} \psi_{-b_1}^* \dots \psi_{-b_{k+1}}^* \psi_{a_{k+1}} \dots \psi_{a_1}$$

where $\lambda_{(d=k+1)}$ now stands for the partition that corresponds to the partition-ordered monomial adjacent to it. This completes the proof of (58).

3.22. **Expanding $(N-1)$ exponentials.** Using (58), we obtain

$$\begin{aligned}
 (68) \quad e^{X_0} \dots e^{X_{N-2}} &= 1 + \sum_{d=1}^{N-1} \sum_{0 \leq a_d < \dots < a_1 \leq N-2} X_{a_d} \dots X_{a_1} \\
 &= 1 + \sum_{d=1}^{N-1} \sum_{\substack{0 \leq a_d < \dots < a_1 \leq (N-2) \\ 1 \leq b_d < \dots < b_1 \leq N}} (-1)^{b_1 + \dots + b_d} d_{\lambda_d} \psi_{-b_1}^* \dots \psi_{-b_d}^* \psi_{a_d} \dots \psi_{a_1}
 \end{aligned}$$

From the remarks following (26), we note that the terms on the right hand side of (68) generate every fermionic expression that corresponds to any partition that fits in the rectangle $[(N-1)^N]$. Additionally, (58) shows that the fermionic expressions are accompanied by the required coefficient with the required sign. Thus

$$\begin{aligned}
 (69) \quad \langle 0 | e^{H(\mathbf{t})} e^{X_0} \dots e^{X_{N-2}} | 0 \rangle \\
 &= d_\phi(\mathbf{v}) \chi_\phi(\mathbf{t}) + \sum_{k=1}^{N-1} \sum_{0 \leq a_k < \dots < a_1 \leq N-2} \sum_{1 \leq b_k < \dots < b_1 \leq N} d_{\lambda_k}(\mathbf{v}) \chi_{\lambda_k}(\mathbf{t}) \\
 &= \sum_{\lambda \subseteq (N-1)^N} d_\lambda(\mathbf{v}) \chi_\lambda(\mathbf{t})
 \end{aligned}$$

Multiplying both sides of (69) by $c_N c_\phi$, we obtain (30).

4. DISCUSSION

The result of this note is that a central object in the algebraic Bethe Ansatz (ABA) approach to the six vertex model (or more precisely the XXZ spin chain), that is the domain wall partition function Z_N , is (a restriction of) a KP τ function, for all values of the crossing parameter. In a sense, what we do is to propose a KP τ function that depends on infinitely many time variables, then restrict the time variables to be power sums of exponentials of the ABA auxiliary space rapidities of Z_N to obtain Lascoux's determinant expression of the latter.

This of course is not the first time that a connection between objects in a solved statistical mechanical model and in a classical integrable hierarchy is observed. For pioneering works in this direction, see [5] and references therein. But in earlier studies that involved the six vertex model or XXZ spin chain, a connection with an integrable nonlinear differential equation (in that case, the nonlinear Schrödinger equation) was made at the free fermion point [5]. The connection with KP that we obtain in this note is valid for all values of the crossing parameter.

We also obtained an expression for Z_N as an expectation value of exponentials of bilinears in the KP fermions. It remains to use this expression to obtain new results, but we observe that the bilinears that appear have a simple combinatorial interpretation. This leads us to believe that analogous expressions can be found for other interesting objects in the ABA approach to the six vertex model and XXZ spin chain such as the scalar products and norms of Bethe vectors.

ACKNOWLEDGEMENTS

We thank Professors M Jimbo, N Kitanine and P Zinn-Justin for useful remarks on the subject of this work and related topics. MW and MZ are supported by Australian Postgraduate Awards and the Department of Mathematics and Statistics, The University of Melbourne.

REFERENCES

- [1] V E Korepin, Commun Math Phys **86** (1982) 391–418.
- [2] A G Izergin, Sov Phys Dokl **32** (1987) 878–879.
- [3] A Lascoux, *Square ice enumeration*, Seminaire Lotharingien de Combinatoire **42**, ‘*The Andrews Festschrift*’, (1999), 30, <http://www.phalanstere.univ-mlv.fr/~al>
- [4] R J Baxter, *Exactly solved models in statistical mechanics*, Second Edition, Dover, 2008.
- [5] V E Korepin, N M Bogoliubov and A G Izergin, *Quantum inverse scattering method and correlation functions*, Cambridge University Press, 1993.
- [6] I G Macdonald, *Symmetric functions and Hall polynomials*, Oxford University Press, 1995.
- [7] T Miwa, M Jimbo and E Date, *Solitons*, Cambridge University Press, (2000).
- [8] R Hirota, *The direct method in soliton theory*, Cambridge University Press, (2004).

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MELBOURNE, PARKVILLE, VICTORIA 3010, AUSTRALIA.

E-mail address: foda, mwheeler, mzup@ms.unimelb.edu.au